



The Life Cycle of Dead Trees

April 12, 2022

John J. Battles, Robert York, and Ariel Roughton




Snag Management in Forestry: Wildlife and Wildfire



Raphael and White. 1984. Use of snags by cavity-nesting birds in the Sierra Nevada. *Wildlife Monograph* 86:3-66.


Hot Drought = 147M Dead Trees




EMC Perspective

1. Habitat value of snags recognized and their retention stipulated in Forest Practice Rules.
2. Snags present a risk to forest operations and wildfire hazard mitigation.
3. Need better understanding of dead wood cycle to inform forest management.

Dead wood cycle



Snags are ephemeral*



Keen 1929, 1955;
Raphael and Morrison 1987, 1993
Battles et al. 2015

*Histogram from Hilger et al. 2012

Decay rates are exponential

HARMON, M. E., CROMACK, K., JR., and SMITH, B. G. 1987. Coarse woody debris in mixed-conifer forests, Sequoia National Park, California. *Can. J. For. Res.* **17**: 1265-1272.




FIG. 1. Mean density of *Abies concolor* boles as a function of time since falling. The density of boles on the ground for more than 24 years was adjusted to reflect volume losses.

Blodgett Forest Research Station



Blodgett Forest Managers

Percy Barr: 1934 – 1955

Rudy Grah: 1958 – 1960

Herb Sampert: 1960 – 1976

Bob Heald: 1976 – 2006


Frieder Schurr: 2006 - 2007

Robert York: 2007 – 2018


Ariel Roughton: 2019 –



Compartment 160: Snag demography study



Compartment 160: 2018-2020



Analytical approach

Strength: 2,600+ trees observed over 37 years


Weakness: Long intervals between inventories; interval censored data

Accelerated Failure Time (AFT) Model

- The effect of a fixed covariate Z is to act multiplicatively on the failure time T or additively on $Y = \log T$.

$$Y = \log T = \alpha + \beta^T Z + \sigma \varepsilon$$


- $\exp(\beta)$: regression parameter which can be interpreted as the ratio of failure time per unit change in covariate.
- AFT model postulates a direct relationship between failure time and covariates.
- “Accelerated failure time model are in many ways more appealing because of their quite direct physical interpretation” – Sir David Cox.



Failure time analysis: Species group


1. Cedar remains standing longest

2. Fir falls the fastest



Failure time analysis: Tree size class


1. Larger trees remain standing longer
2. Smallest trees fall fastest
(sub < 7.9 in DBH)




Failure time analysis: Decay class

Stubs remain standing a long time

Nest trees tend to fall the fastest



Median Fall Rates




Weibull additive model with species group and size class was the best model

| Species | DBH | Fall Times (years) | | |
|---------|-------|--------------------|------|------|
| | | median | 25th | 75th |
| CEDAR | sub | 17.9 | 11.1 | 26.0 |
| CEDAR | small | 19.1 | 11.9 | 27.8 |
| CEDAR | med | 18.4 | 11.4 | 26.7 |
| CEDAR | large | 17.2 | 10.7 | 24.9 |
| PINE | sub | 16.4 | 10.2 | 23.9 |
| PINE | small | 12.7 | 7.9 | 18.5 |
| PINE | med | 13.5 | 8.4 | 19.7 |
| PINE | large | 18.6 | 11.6 | 27.1 |
| OAK | sub | 1.1 | 0.7 | 1.7 |
| OAK | small | 14.8 | 9.2 | 21.5 |
| OAK | med | 14.2 | 8.8 | 20.7 |
| OAK | large | 15.5 | 9.6 | 22.5 |
| FIR | sub | 8.2 | 5.1 | 12.0 |
| FIR | small | 13.8 | 8.6 | 20.0 |
| FIR | med | 13.6 | 8.5 | 19.7 |
| FIR | large | 14.4 | 9.0 | 21.0 |

Decay progression

(wildlife snags only)

| Decay Class | Longevity (years) | |
|-------------|-------------------|-----|
| | mean | se |
| 1 | 2.0 | 1.4 |
| 2 | 2.1 | 1.3 |
| 3 & 4 | 6.8 | 4.8 |
| 5 | 9.3 | 6.2 |



EMC Application

Snags typically remain standing for 15 years.

Only 9 years of their 15 year median “lifetime” do snags provide critical wildlife habitat.

At Blodgett, it takes a minimum of 20 years to grow a tree big enough to provide wildlife habitat.

Current snag retention guidelines do not consider longevity of existing snags.

Guidelines to not include provisions for snag recruitment in harvested stands.

Results suggest that one matching live tree should be retained for every snag retained to “recruit” replacements for snags that fall.

Decay rate of coarse woody debris

Log cemeteries established in 2018

Initial samples collected

Initial samples processed for wood density

2019: Monumented logs for the long-term





Department of Environmental Science, Policy, &
Management

Retrospective Analysis of Wood Decay by Species in Sierran Mixed Conifer

By Adam Sawicky



Mentor: Dr. John J. Battles

Approach

Conduct a field inventory of downed logs


Match records of fall; Evaluate current decay status

Use paired information to estimate decay rate




Rates of decay

| Species Group | n | Decay Rate (g/cm ³ yr ⁻¹) |
|---------------|----|---|
| Softwood | 62 | 0.013 (0.002) |
| Cedar | 60 | 0.007 (0.002) |
| Hardwood | 28 | 0.017 (0.004) |



Decay rates at Blodgett: Slower than expected



Goal: Improve dead wood cycle in forest growth models

Existing models:

1. Underestimate snag fall rates
2. Overestimate CWD decay rates

Major Challenge: Generalize

